

Frontier Analysis, Ltd

TECHNICAL SERVICE RESPONSE NO.: UT042

Subject: Analysis of Black Material on the Roots and Inside Corn Stalks from a Crop Formation (Mission, B.C., August 2002)

Date: May 25, 2005

Requested By: Nancy Talbott
BLT Research

Reported By: P. A. Budinger
Analytical Scientist

Background/Objective:

The background as described by Nancy Talbott follows.

“The enclosed corn stalks come from a crop formation in Mission B.C., Canada which occurred in late August, 2002. You’ll note that the stalks have a lot of black on them. The stuff on the leaves I suspect is from fungus. But there is a different quality to the black on the roots and inside the corn stalks near the bottom which we are wondering about. Could this be a carbon blackening? Like perhaps one might see in a lightning strike? Or is it just the fungus?”

The objective is to determine the cause of the black color on the ends and roots of the corn stalks.

An aerial photograph of the site of the corn formation is on the following page.




*Corn stalks taken from large circle (#2)
on Oct. 13, 2002*

Conclusions:

- The analysis shows the black interior is corn fungus (ustilago). No carbon, which would be indicative of burning/heat, was detected.
- Two tests for determining the difference between carbon and black corn fungus can be easily done by field workers. The first would be to smell for charred material. The second would involve touching the black area. If the black material is carbon, some will come off and blacken the fingers. Corn fungus would not come off.

Procedure:

The samples consist of four corn stalks identified as “stalks from crop circle “B” by river”. Following is a diagram of the formation, which shows the two sampling locations of the stalks.



Mission, British Columbia #2
Late August, 2002


Lay and Measurements Diagram
(Field Survey: September 22, 2002)

© Paul Anderson, CCCRN

Infrared spectra were obtained from the black areas of the ends and the roots of the stalks, as well as a ‘clean’ area inside of the stalk for reference. Additionally, a reference spectrum was obtained from corn fungus (ustilago) on a corn husk from Budinger’s compost pile. The data were obtained a Nicolet Avatar 360 spectrometer using the Harrick SplitPea™ sampling accessory. Photographs were taken with a Kodak Digital Science MDS 120 camera.

Results:

Photographs of the stalks follow.



The Stalks



Blackened Ends



End (Close Up)




Blackened Roots

On receipt of the sample the black ends and root areas were touched to determine whether any carbon would be transferred to the fingers. None was. The blackened areas were also smelled for the presence of charred material. Again no characteristic odor was detected. This was the first clue that the black material may not be carbon. This was rather surprising to this analyst due to the intense blackness of these areas.

Infrared spectra of the blackened ends and root areas are identical to each other and match a reference of corn fungus (ustilago). No carbon was detected. If


carbon were present, the spectra would have displayed significant light scattering in the higher frequency regions ($4000 - 1800 \text{ cm}^{-1}$). Following are infrared spectra of the black areas of an end and the root which are compared to a reference of corn fungus. Clearly all spectra are identical.

Infrared Spectra of Corn Stalk Black End and Black Root and a Corn Fungus Reference



And additional note should be made regarding the spectrum of ustilago compared to a 'clean' area of the interior of the corn stalk for the scientist familiar with infrared interpretation. The spectra appear to be similar, however, there are significant differences. The ustilago spectrum is clearly typical of fungus. The ustilago fungus spectrum shows: increase in C-H stretch absorption ($2000-2800 \text{ cm}^{-1}$); additional ester C=O (1740 cm^{-1}); increased absorption contributed to by an amide C=O ($1700-1490 \text{ cm}^{-1}$); increased absorption to a combination of modes ($1480-1290 \text{ cm}^{-1}$); loss of resolution in the C-O stretch region ($1140-930 \text{ cm}^{-1}$). It is very close to a reference of dried wood ear mushroom which is also a fungus. Following are spectra of the clean corn stalk interior, corn fungus (ustilago), and dried wood ear mushroom illustrating the above.

Infrared Spectra of 'Clean' Corn Stalk Interior, Dried Wood Ear Mushroom, and Corn Fungus (*Ustilago*)



File: UT042

Phyllis A. Budinger